

## Preliminary studies on *in vitro* methods for the evaluation of anticoccidial efficacy/resistance in ruminants



Ane Odden<sup>a,\*</sup>, Snorre Stuen<sup>a</sup>, Heidi L. Enemark<sup>b</sup>, Lucy J. Robertson<sup>c</sup>, José Manuel Molina<sup>d</sup>, Antonio Ruiz<sup>d</sup>

<sup>a</sup> Faculty of Veterinary Medicine, Department of Production Animal Clinical Sciences, Norwegian University of Life Sciences, Kyrkjavegen 332/334, N-4325, Sandnes, Norway

<sup>b</sup> Norwegian Veterinary Institute, Ullevålsveien 68, P.O. Box 750 Sentrum, N-0106, Oslo, Norway

<sup>c</sup> Faculty of Veterinary Medicine, Department of Food Safety and Infection Biology, Norwegian University of Life Sciences, P.O. Box 8146 Dep, N-0033, Oslo, Norway

<sup>d</sup> Parasitology Unit, Department of Animal Pathology, Faculty of Veterinary Medicine, University of Las Palmas de Gran Canaria, 35416, Arucas, Las Palmas, Spain

### ABSTRACT

#### Keywords:

*Eimeria* spp.

Anticoccidial resistance

*In vitro* assay

Bovine colonic epithelial cells

Ovine *Eimeria* spp. infections cause increased mortality, reduced welfare and substantial economic losses, and anticoccidials are important for their control. Recent reports of anticoccidial resistance against ovine *Eimeria* spp. necessitate the development of *in vitro* methods for the detection of reduced anticoccidial efficacy, especially since the *in vivo* methods are both expensive, time consuming and requires the use of otherwise healthy animals. The aim of the present study was therefore to approach a preliminary standardization of *in vitro* assays for evaluation of the efficacy of the most commonly used anticoccidials in ruminants. For this purpose, apart from the evaluation of inhibition of oocyst sporulation, most effort was concentrated on assessment of the capacity of the different anticoccidials to inhibit both the invasion and further development (up to the first schizogony) of *E. ninakohlyakimovae* sporozoites in bovine colonic epithelial cells (BCEC). For this purpose, infected cultures were monitored 1, 8 and 15 days post infection to determine the infection rate, number of immature schizonts and number, size and appearance of mature schizonts, respectively. No clear inhibitory effect was found with any of the anticoccidial formulations tested, and we could not identify why there were no measurable effects from the different anticoccidials. Despite the lack of positive results, further investigations should be encouraged, as this could decrease the need for animal experiments and could be used in the initial assessment of anticoccidial efficacy of new drugs.

### 1. Introduction

Infections caused by *Eimeria* species are some of the most important parasitic diseases affecting the profitability of ruminant production systems (Keeton and Navarre, 2018). Young animals are particularly affected by clinical disease, often in the period around weaning. Infection may result in diarrhoea, reduced growth and occasional deaths (Dauguschies and Najdrowski, 2005; Ruiz et al., 2006; Chartier and Paraud, 2012). The control of ruminant coccidiosis is traditionally based on the combination of good management together with prophylactic or metaphylactic treatment with anticoccidials (Dauguschies and Najdrowski, 2015). However, for at least a decade no new drugs have been brought to market despite evidence of anticoccidial resistance

(ACR) to current treatments occurring worldwide in poultry (McDougald et al., 1987; Peek and Landman, 2005; Lan et al., 2017). Toltrazuril resistance has been confirmed in porcine *Cystoisospora suis* (Sheresta et al., 2017), and in ovine *Eimeria* spp. (Odden et al., 2018a). Toltrazuril resistance in ovine *Eimeria* spp. in Norway is probably related to the widespread and extensive use of anticoccidial (AC) treatment in this country, as discussed in a recent publication based on a questionnaire study and aiming to identify potential risk behaviour for development of ACR (Odden et al., 2017).

Standard methods for *in vivo* evaluation of anthelmintic efficacy are not valid for the assessment of the efficacy of anticoccidial drugs, due to the substantial lifecycle differences between nematodes and coccidia. Accordingly, Odden et al. (2018b) recently published a new

**Abbreviations:** ACE, anticoccidial efficacy; ACR, anticoccidial resistance; BCEC, bovine colonic epithelial cell; BUVEC, bovine umbilical vein endothelial cells; CUVEC, caprine umbilical vein endothelial cells; FECRT, faecal egg count reduction test; FOCRT, faecal oocyst count reduction test; MIC, minimal inhibitory concentration; MDBK, Madin-Darby bovine kidney cells; WAAVP, Word Association for Advanced Veterinary Parasitology

\* Corresponding author.

E-mail address: [ane.odden@nmbu.no](mailto:ane.odden@nmbu.no) (A. Odden).

<https://doi.org/10.1016/j.exppara.2019.04.009>

Received 20 January 2019; Received in revised form 8 April 2019; Accepted 24 April 2019

Available online 26 April 2019

0014-4894/© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

approach for field evaluation of anticoccidial efficacy (ACE) against ovine *Eimeria* spp. using a method based on the WAAVP recommended faecal egg count reduction test (FECRT) for identifying resistance to anthelmintics (Coles et al., 1992), with certain modifications. Changes from the original protocol included the use of geometric, instead of arithmetic, means and restriction to the exponential phase of oocyst excretion for a more accurate evaluation of the ACE. The usefulness of this new approach, named FOCRT (Faecal Oocyst Count Reduction Test), as a tool to evaluate ACE in the field was shown by the same authors in a controlled efficacy study in which the existence of reduced efficacy of toltrazuril in a field isolate of ovine *Eimeria* spp. was demonstrated (Odden et al., 2018a). In that study, 50% of the experimentally *Eimeria*-infected lambs were metaphylactically treated with the recommended dose of 20 mg/kg toltrazuril (Baycox® Sheep vet., Bayer Animal Health), but no difference in oocyst excretion between treated and control lambs was observed (Odden et al., 2018a). Furthermore, there were no differences in weight gain and macro-/microscopic findings at post-mortem examination.

The *in vitro* evaluation of drug efficacy or resistance against helminths has been extensively documented. As reviewed by Taylor et al. (2002), the available *in vitro* tests are diverse, and include: egg hatch assays, migration and motility assays, and larval and adult development tests, as well as biochemical and molecular techniques. Most of these *in vitro* tests have been used to detect and describe resistance against most anthelmintic groups, including benzimidazoles (Rialch et al., 2013; Ramünke et al., 2016; Milhes et al., 2017), imidazothiazoles (Martines-Valladares et al., 2013), and macrocyclic lactones (Almeida et al., 2013; Milhes et al., 2017). The same methodology has also been applied for the evaluation of the efficacy of a number of biological compounds or plant extracts in the last decades (Alawa et al., 2003; Iqbal et al., 2006; Al-Rofaai et al., 2012; Araújo et al., 2017; Jaso Díaz et al., 2017; Novobilsky et al., 2013). However, as for *in vivo* assessments, most of these *in vitro* tests are not appropriate for evaluation of ACE due to the complexity and particularities of the endogenous and exogenous life-cycle of *Eimeria* spp.. Accordingly, new *in vitro* methods have recently been developed, mostly for investigation of poultry coccidiosis. For instance, there are *in vitro* assays showing that different plant extracts inhibit the sporulation of oocysts, the viability of sporozoites, or the invasion rate of different *Eimeria* species of poultry (Molan et al., 2009; Khalafalla et al., 2011; Burt et al., 2013; Gadelhaq et al., 2018). Further evaluation of the efficacy of different anticoccidials has been accomplished by using a combination of cell culture and qPCR (Thabet et al., 2015). Minimum inhibitory concentrations (MIC) of monensin, maduramicin, salinomycin and lasalocid have been determined, based on the development to mature schizonts in Madin-Darby bovine kidney (MDBK) cells of sporozoites of a field *Eimeria tenella* strain (Thabet et al., 2015). The same authors recently proposed *in vitro* *E. tenella* assays as a replacement for animal experiments for ACE testing (Thabet et al., 2017). The efficacy of the same polyether ionophores described above and toltrazuril were tested (Thabet et al., 2017) using *in vitro* sporozoite inhibition and reproduction inhibition assays and further determination of the MIC.

In contrast to poultry, limited information is available on *in vitro* assays evaluating the ACE for ruminant *Eimeria* species. Interestingly, the launch of a *Guideline for Evaluating the Efficacy of Anticoccidials in Mammals* (Joachim et al., 2018), raised concerns about drug-resistance testing and alternative methods for evaluation of drug efficacy. For both *in vivo* and *in vitro* tests, the authors stress the need for defined strains of ruminant *Eimeria* species for protocol standardization. Ruiz et al. (2013a) reported the isolation and experimental infection of a defined strain of *E. ninakohlyakimovae*, initially isolated from the field in 2006 in Gran Canaria (Spain). This *Eimeria* strain has been subsequently used in a number of immunological, pathological, and immunoprophylactic studies (Ruiz et al., 2013b, 2014; Pérez et al., 2015, 2016; Matos et al., 2017a, 2017b, 2018).

Against this background, the aim of the present study was to address

preliminary standardization of *in vitro* assays for evaluation of the efficacy of the most commonly used anticoccidials in ruminants. For this purpose, apart from the evaluation of inhibition of oocyst sporulation, most effort was concentrated on assessment of the capacity of the different anticoccidials to inhibit both the invasion and further development (up to the first schizogony) of *E. ninakohlyakimovae* sporozoites in bovine colonic epithelial cells (BCEC).

## 2. Material and methods

### 2.1. Parasite maintenance

The *E. ninakohlyakimovae* strain, GC, used in the present study was initially isolated from goats in the Gran Canaria Islands (Spain), and maintained by passage in goat kids for oocyst production. Oocysts were isolated according to Jackson (1964) with some modification. Briefly, faeces were mixed 1:1 with water and passed through sieves of decreasing pore diameter, down to approximately 100 µm. The faecal mix was subsequently mixed 1:1 with saturated sugar-solution and floated onto glass slides, which were washed every 2 h with distilled water, for three consecutive days. The washings were centrifuged at 2300 × g for 20 min, the supernatant was discarded, and the resulting sediment mixed 1:1 with distilled water in a glass flask. Oocysts collected in the flask were set to sporulate under constant aeration for 7 days at room temperature (RT). Sporulated oocysts were stored at 4 °C in culture flasks (Nunc) with access to air.

Oocyst purification and isolation of sporozoites were performed according to Fayer and Hammond (1967) and Pérez et al. (2015), with slight modifications. Sporulated oocysts were added to 5% (w/v) sodium hypochlorite and stirred on ice for 30 min using a magnetic flea, followed by centrifugation at 233 × g, 10 °C for 5 min. The supernatant was mixed 1:1 with distilled water and centrifuged at 1500 × g, RT for 10 min. The resulting sediment, containing oocysts, was suspended in sterile 0.5% L-cysteine (C<sub>2</sub>H<sub>7</sub>NO<sub>2</sub>S, Merck) and 1.68% NaHCO<sub>3</sub> solution (Sigma Aldrich) and incubated at 100% CO<sub>2</sub> atmosphere, 37 °C for 20 h. Subsequently, oocysts were suspended in Hank's balanced salt solution (HBSS, Sigma Aldrich) containing 0.075% w/v % trypsin (Biowest), 0.15% w/v sodium taurodeoxycholate (Sigma Aldrich), and 8% sterile-filtered bovine serum from two different animals obtained from the local abattoir. The oocyst suspension was incubated and checked periodically (approximately every 30 min) by microscopy for up to 4 h (37 °C, 5% CO<sub>2</sub> atmosphere). Excysted sporozoites were washed three times (20 min, RT, 1500 × g) and suspended in RPMI-1640 (Sigma-Aldrich) medium with the appropriate anticoccidial concentration (see Table 1) and transferred to the cell cultures.

### 2.2. Cell culture

Bovine colonic epithelial cells (BCEC) (Ruiz et al., 2010) were cultured in RPMI-1640 medium (Sigma-Aldrich) in 12-well plates and incubated at 37 °C and a 5% CO<sub>2</sub> atmosphere until confluence. The

**Table 1**

Concentrations (µg/ml) of the different commercial anticoccidials included in the assay.

	Concentration	Toltrazuril	Diclazuril	Decoquinatate	Sulphonamide	Control DMSO
		T	D	Q	N	X
A	25.0	25.0			25.0	25.0
B	15.0	15.0			15.0	15.0
C	10.0	10.0	1.0		10.0	10.0
D	5.0	5.0	0.1		5.0	5.0
E	1.0	1.0	0.01		1.0	1.0
F	0.1	0.1	0.001		0.1	0.1
G	0.01	0.01	0.0001		0.01	0.01
H	0.001	0.001	0.00001		0.001	0.001

medium was supplemented with 500 U/ml penicillin (Sigma-Aldrich), 50 µg/ml streptomycin (Sigma Aldrich), 0.25 µg/ml amphotericine (Sigma Aldrich), 0.5 µg/ml Plasmocin™ (InvivoGen), and 20% faecal calf serum (Biowest).

### 2.3. Anticoccidials

Both commercial anticoccidial formulations (toltrazuril (Baycox® Sheep vet., Bayer Animal Health), diclazuril (Rumicox®, Steve Veteriaria), decoquinat (Deccox®, Zoetis) and sulphonamide (Cunitotal®, Ecuphar)), and the source anticoccidial toltrazuril and its metabolites, toltrazuril sulphoxide and toltrazuril sulphone (Sigma Aldrich), were used in the study. Additionally, two different negative controls, dimethyl sulphoxide (DMSO, Sigma Aldrich) for the commercial anticoccidials and dimethyl formamide (DMF, Sigma Aldrich) for the pure source/metabolite anticoccidials, were also included in the assays. The anticoccidials were diluted up to a 100 µg/ml stock concentration, either by using DMSO (commercial anticoccidials) or DMF (pure source/metabolite anticoccidials); further dilutions were made using cell culture medium. The anticoccidial concentrations applied to sporozoites and cells are shown in Table 1 (commercial anticoccidials) and Table 2 (pure source/metabolite anticoccidials). The DMSO and DMF maximum concentrations used were 0.4%. Anticoccidials were continuously included in the cell-culture medium throughout the study under the assumption that the drug concentrations remained stable. Monensine (Sigma Aldrich), at the same concentrations used for pure/derivate anticoccidials, was employed as positive control.

### 2.4. Infection of host cells and evaluation of invasion and development

Confluent BCEC cell layers containing anticoccidials or negative controls were infected with 100,000 freshly excysted sporozoites and incubated at 37 °C and 5% CO<sub>2</sub>, in duplicates. The culture medium was changed 24 h after infection and subsequently three times a week.

At 24-h post infection, the invasion rate was determined by examining 15–20 photos taken at 400× magnification using a phase-contrast microscope (DMIL, Leica) and associated digital camera DFC299. Photos were taken systematically, covering the central area of the wells with the highest phase contrast, and scanning in a zig-zag motion. The scanning movement meant that a single field was not photographed more than once. For each of the 15–20 pictures taken of a single culture (condition), the number of intracellular sporozoites was enumerated using Carl Zeiss ZEN 2.3 lite software. To determine the average number of cells per picture, 600 pictures were evaluated. Invasion rate is given as the percentage of infected cells 24-h post infection.

Evaluation of development was assessed at days 8 and 15 post infection. At day 8, 15–20 pictures were taken systematically at 400× magnification as described for day 1, and immature schizonts were counted. Pictures were only taken when at least one immature schizont was present in a field, and the total number of fields examined, with or without immature schizonts, up to a maximum of 50 fields, was recorded. This early development is presented as the number of immature

**Table 2**

Concentrations (µg/ml) of the pure source/metabolite anticoccidials included in the assay.

Concentration	Pure toltrazuril	Toltrazuril sulfoxide	Toltrazuril sulfone	Control DMF
	TT	TD	TN	DMF
A	25.0	25.0	25.0	25.0
B	5.0	5.0	5.0	5.0
C	1.0	1.0	1.0	1.0
D	0.1	0.1	0.1	0.1
E	0.01	0.01	0.01	0.01

schizonts per mm<sup>2</sup>.

At day 15, the mature schizonts were counted at 100× magnification over 15–20 pictures taken as referred previously for each condition. The degree of development is given as the number of schizonts per mm<sup>2</sup>. In addition, the size and appearance of the schizonts were described.

### 2.5. In vitro oocyst sporulation inhibition assay

The effects of the different anticoccidials on sporulation of *E. nakholyakimovae* oocysts *in vitro* were evaluated as follows. Briefly, 1.5 ml Eppendorf tubes were filled with a 20 µl suspension of 5000 unsporulated oocysts in sterile PBS, 30 µl distilled H<sub>2</sub>O, 50 µl of 10% potassium dichromate, and 100 µl of different anticoccidials (A-D: 0.5, 0.1, 0.05 and 0.025 mg/ml). The incubations were performed at RT for 24 h and then the oocysts were washed in distilled water four times (2000×g, 10 min). After the last washing step, the oocysts were re-suspended in 1 ml 2% potassium dichromate, transferred into 24-multiwell tissue culture plates (Nunc) and incubated in the presence of oxygen at RT. As negative controls, similar concentrations of DMSO (commercial anticoccidials) or DMF (pure source/metabolite anticoccidials) to those used to dissolve the corresponding anticoccidials (from 8% to 1%) were used, and serial formaldehyde solutions (Panreac) served as positive controls (4, 2, 1 and 0.5%). Oocyst sporulation rate was determined after 72 h at RT by microscopic analysis, and for this purpose a minimum of 100 oocysts were counted and analysed. The assay was repeated twice including duplicates of all concentrations in both assays.

### 2.6. Statistical methods


Data were managed and analysed in Excel (2013) (Microsoft Inc.), in addition to analyses performed in R (R Core Team, 2017) and Sigmaplot 12.0. Data from the different *in vitro* analysis (invasion rate, development of immature and mature schizonts and inhibition of sporulation) were grouped and analysed independently for significant differences by using Chi-square tests. Effects of the different anticoccidial concentrations on the parasites were compared to the corresponding DMSO or DMF negative controls. Differences were regarded as significant at a level of P < 0.05.

## 3. Results

### 3.1. Sporozoite invasion

The average sporozoite invasion rate ranged from a minimum of 14.1% infected cells (25.0 µg/ml toltrazuril) to a maximum of 39.9% (control DMSO). The highest invasion rate for the commercial anticoccidials was seen for 0.1 µg/ml decoquinat at 39.7%, and for the pure source/metabolite anticoccidials for 5 µg/ml TN at 39.3%. The lowest invasion rate was seen for 25.0 µg/ml toltrazuril at 14.1% and for 0.1 µg/ml toltrazuril sulphoxide TD at 20.2%, for the commercial and pure source/metabolite anticoccidials, respectively. The average invasion rates for the controls, combining all concentrations, were 33.8 ± 1.1% (range: 26.6–39.9%) and 29.9 ± 1.6% (range: 27.4–33.2%) for DMSO and DMF, respectively.

The sporozoite invasion showed a significant difference (P < 0.05) for toltrazuril (25.0 µg/ml) and diclazuril (25.0 µg/ml), compared with the corresponding control DMSO concentration, but not for any of the other anticoccidials, or concentrations (Fig. 1). Although only the highest concentration of commercial toltrazuril was significantly different from the control, the following three concentrations showed a gradual increase in the percentage of infected cells of 22.6, 31.5, and 34.1%, respectively.



**Fig. 1.** Invasion rates of *Eimeria ninkohlyakimovae* sporozoites into bovine colonic epithelial cells under the influence of commercial (A) or pure source/metabolite (B) anticoccidials, calculated 24 h post infection. A: (T) toltrazuril, (D) diclazuril, (Q), decoquinat, (N) sulphonamide, (X) dimethyl sulphoxide – negative control. B: (TT) pure toltrazuril, (TD) toltrazuril sulphoxide, (TN) toltrazuril sulphone, (DMF) dimethyl formamide negative control. See Tables 1 and 2 for specific concentrations (A–H) used for each anticoccidial.

### 3.2. Sporozoite development


#### 3.2.1. Immature schizonts

The average number of immature schizonts ranged from a minimum of 4.7 schizonts/mm<sup>2</sup> (25.0 µg/ml toltrazuril) to a maximum of 46.3 schizonts/mm<sup>2</sup> (5.0 µg/ml control DMF). The average number of schizonts/mm<sup>2</sup> was of 22.2 ± 0.8 and 36.9 ± 2.0 for DMSO and DMF, respectively. The highest number of schizonts/mm<sup>2</sup> recorded for the commercial anticoccidials was 125, found with two different drugs; 0.1 µg/ml toltrazuril and 1.0 µg/ml diclazuril, whereas the highest number recorded for the pure source/metabolite anticoccidials was 204.5 schizonts/mm<sup>2</sup> (1.0 µg/ml toltrazuril sulphone).

There was a significant difference ( $P < 0.05$ ) in sporozoite development into immature schizonts at day 8 for 25.0 µg/ml of toltrazuril and diclazuril, compared with their corresponding concentration of the control DMSO (Fig. 2). This difference was not found in any of the other anticoccidials, at any concentrations.

#### 3.2.2. Mature schizonts

The average number of mature schizonts/mm<sup>2</sup> ranged from a minimum of 3.2 ± 0.6 schizonts/mm<sup>2</sup> (sulphonamide 25 µg/ml) to a maximum of 21.6 ± 1.6 schizonts/mm<sup>2</sup> (DMSO) (Fig. 3). The lowest number of schizonts/mm<sup>2</sup> recorded for the commercial anticoccidials was 2.7 (Sulphonamide 25 µg/ml), whereas for the pure source/



**Fig. 2.** Development of immature *Eimeria ninkohlyakimovae* schizonts in bovine colonic epithelial cells under the influence of commercial (A) or pure source/metabolite (B) anticoccidials evaluated 8 days post infection. A: (T) toltrazuril, (D) diclazuril, (Q), decoquinat, (N) sulphonamide, (X) dimethyl sulphoxide – negative control. B: (TT) pure toltrazuril, (TD) toltrazuril sulphoxide, (TN) toltrazuril sulphone, (DMF) dimethyl formamide negative control.


metabolite anticoccidials the lowest number recorded was 5.3 schizonts/mm<sup>2</sup> (toltrazuril, 5 mg/ml). For commercial anticoccidials, in particular, the schizont numbers at the highest concentration were lower than for negative controls but significant differences could not be demonstrated. The same was observed for the mean appearance of the mature schizonts at day 15 after infection, whose values ranged from 1.3 to 2.0 for all anticoccidials and corresponding controls (Fig. 4). In contrast, the size of the mature schizonts remained relatively similar, with diameters fluctuating between 46.2 ± 6.9 µm (decoquinat, 10 µg/ml) and 76.7 ± 7.1 (toltrazuril 25 µg/ml) (Fig. 5). Schizonts ≥ 50 µm amounted for 83.2% of all schizonts counted. However, by only evaluating large schizonts (≥ 50 µm), the results did not significantly change, from the evaluations of all schizonts.

### 3.3. Oocyst sporulation inhibition assay (OSIA)

None of the anticoccidials, commercial or pure source/metabolite, appeared to influence oocyst sporulation rate, which ranged from 81.3% (pure toltrazuril, 0.5 mg/ml) to 94.6% (diclazuril, 0.1 mg/ml). In contrast, less than 30% oocysts sporulated after incubation with the positive control (formaldehyde; data not shown).

## 4. Discussion


In the present study we assessed the ability of *E. ninkohlyakimovae*



**Fig. 3.** Number of mature *Eimeria ninkohlyakimovae* schizonts in bovine colonic epithelial cells under the influence of commercial (A) or pure source/metabolite (B) anticoccidials evaluated 15 days post infection. A: (T) toltrazuril, (D) diclazuril, (Q), decoquinatate, (N) sulphonamide, (X) dimethyl sulphoxide – negative control B: (TT) pure toltrazuril, (TN) toltrazuril sulphone, (TD) toltrazuril sulphoxide, (DMF) dimethyl formamide – negative control. The data are represented as the mean duplicates  $\pm$  STD. See Tables 1 and 2 for specific concentrations (A–H) used for each anticoccidial.

sporozoites to invade and further develop up to first schizogony in BCECs in the presence of different concentrations of anticoccidials or controls. No clear inhibitory effect was found with any of the anticoccidial formulations tested. Similar to our results, Thabet et al. (2017) did not find a significant correlation between *in vivo* data and percentage of reproduction inhibition *in vitro* for toltrazuril when evaluating the development of *Eimeria tenella* in MDBK cells. In addition, compared with different ionophores, toltrazuril showed the highest value of minimum inhibitory concentration. The authors suggested that those data probably indicated that this test system is not appropriate for assessing toltrazuril sensitivity of *E. tenella*, and the same may be the case for the non-ionophore anticoccidials evaluated here.

The commercial anticoccidials, toltrazuril (Baycox<sup>®</sup>), diclazuril (Rumicox<sup>®</sup>), and decoquinatate (Deccoq<sup>®</sup>), assessed in the present study have been demonstrated to be effective at reducing parasite burdens and have been associated with increased growth rates in calves (Dauguschies and Najdrowski, 2005; Mundt et al., 2005; Enemark et al., 2015), lambs (Taylor et al., 2011; Taylor and Bartran, 2012; Diaferia et al., 2013), and goat kids (Foreyt et al., 1986; Ruiz et al., 2012; Iqbal et al., 2013); all three anticoccidials are registered in different EU countries for cattle and sheep. In contrast, in many countries sulphonamides are no longer authorized for anticoccidial treatment, although some derived drugs, such as sulphadimethoxine, are still commercially




**Fig. 4.** Size of mature *Eimeria ninkohlyakimovae* schizonts in bovine colonic epithelial cells under the influence of commercial (A) or pure source/metabolite (B) anticoccidials evaluated 15 days post infection. A: (T) toltrazuril, (D) diclazuril, (Q), decoquinatate, (N) sulphonamide, (X) dimethyl sulphoxide - negative control, B: (TT) pure toltrazuril, (TN) toltrazuril sulphone, (TD) toltrazuril sulphoxide, (DMF) dimethyl formamide – negative control. The data are represented as the mean duplicates  $\pm$  STD. See Tables 1 and 2 for specific concentrations (A–H) used for each anticoccidial.

available for cattle coccidiosis, e.g., in the USA (Burke et al., 2013).

Despite the strong efficacy documented *in vivo* for the four commercial anticoccidials evaluated in our study, limited evidence of their activity *in vitro* could be demonstrated in the cell-culture model employed here. Indeed, only the highest concentration of toltrazuril (Baycox<sup>®</sup>) and diclazuril (Rumicox<sup>®</sup>) significantly reduced the infection rate as well as the number of immature schizonts. Low numbers of mature schizonts were also found for the two higher concentrations of both commercial anticoccidials, but the differences were not statistically significant; this probably reflects the relatively high standard deviations associated with a limited number of replicates.

Bioconversion of antiparasitic drugs in the host is not uncommon (Lanusse et al., 1995), so the apparent lack of effect of the commercial anticoccidials analysed here could be related to the inability of the current *in vitro* system to metabolize the drugs to active compounds. To assess this possibility, the anticoccidial activity *in vitro* of pure toltrazuril (without excipient) and two of its main metabolites were evaluated at different concentrations. Within the treated host, toltrazuril undergoes extensive metabolism to toltrazuril sulphoxide and then to toltrazuril sulphone (ponazuril) (Lim et al., 2010), which appears to have anticoccidial activity against *Cystoisospora suis* (Bach et al., 2003) and goat *Eimeria* infections (Gibbons et al., 2016). However, far from increasing their anticoccidial effect *in vitro* in the present



**Fig. 5.** Appearance of mature *Eimeria ninakholyakimovae* schizonts in bovine colonic epithelial cells under the influence of commercial (A) or pure/derivate (B) anticoccidials evaluated 15 days post infection. The schizont condition was scored as: 1) healthy and with grey scale colour at phase contrast; 2) relatively healthy and with brownish appearance at phase contrast; 3) unhealthy and with irregular shape. A: (T) toltrazuril, (D) diclazuril, (Q), decoquinat, (N) sulphonomide, (X) dimethyl sulphoxide - negative control, B: (TT) pure toltrazuril, (TN) toltrazuril sulphone, (TD) toltrazuril sulphoxide, (DMF) dimethyl formamide - negative control. The data are represented as the mean duplicates  $\pm$  STD. See Tables 1 and 2 for specific concentrations (A–H) used for each anticoccidial.

study, the metabolites of pure toltrazuril, toltrazuril sulphone and toltrazuril sulphoxide, only showed a minor inhibitory activity against *E. ninakholyakimovae* sporozoite invasion and further development.

As the *in vitro* anticoccidial effects of the drugs analysed here were mostly found at the highest concentrations, a more pronounced inhibitory response would have been expected by increasing the amount of drug available in the cultures. However, concentrations higher than those shown in Tables 1 and 2 could not be estimated in the culture system employed here, due to evidence of cell damage, either by the effect of the drug itself or the concentration of solvent used. In particular, high concentrations of decoquinat (Deccox<sup>®</sup>) showed a strong cytotoxicity, and therefore the two highest concentrations were not evaluated for this commercial anticoccidial. Similarly, the final DMSO percentage for the highest concentration led to a significant reduction in the number of immature schizonts.

Irrespective of the drug metabolite or concentration used, the type of cells might be another important factor to be considered in an *in vitro* system for the evaluation of ACE. The cell line used here, BCEC, has previously been shown to be a suitable *in vitro* model for the development of *E. ninakholyakimovae* up to the first schizogony (Ruiz et al.,

2010), which was also achieved in the present study. Example photographs of intracellular sporozoites, immature schizonts, and mature schizonts containing merozoites are provided in electronic supplementary material. These types of cells are colonic in origin and it is possible that absorption of the anticoccidials used in our study differs between intestinal segments; this is relevant because the pathogenic species mainly infect the small intestine and the caecum (Deplazes et al., 2016). It may also be that the process of creating a permanent cell line (Föllmann et al., 2000), might have influenced the cells' ability to incorporate different substances. Multi-drug resistance has been reported in colonic cancer cells, which block drug activity by efflux transporters that promote metabolism, elimination, and detoxification (Chen et al., 2012); whether the colonic cell line used in our studies share some of these features cannot be excluded and should be evaluated. The use of a different cell line from closely related hosts, possibly of small intestinal origin or primary endothelial origin, like the bovine or caprine umbilical vein endothelial cells (BUVEC and CUVEC, respectively), might thus have provided more useful results. Infections of BUVEC and CUVEC by *E. ninakholyakimovae* sporozoites have also been shown to result in merogony and macromeront-I formation (Ruiz et al., 2010).

The development of an *in vitro* model for the study of biological processes does not necessarily take into account the complexity of circumstances occurring *in vivo*. When evaluating drug efficacy *in vitro*, the assumption is made that the mechanism of action also occurs in the test system, but this may not always be the case. Indeed, the mechanisms of action of the different anticoccidials are not always well elucidated. For example, the proposed mode of action of toltrazuril is thought to be directed against the first and second generation schizonts, microgamonts, and macrogamonts (Mehlhorn, 2008). The action is probably achieved by inhibiting mitochondrial respiration and nuclear pyrimidine synthesis in the parasite, possibly by inhibiting dihydroorotate dehydrogenase (Harder and Harberkorn, 1989). However, the distribution of this enzyme in different intestinal segments is still unknown and molecular-related reactions deserve further investigation. For diclazuril, the mechanisms of action are unknown, but it has been shown that the activity is only directed against specific endogenous stages of *Eimeria* spp. (Mehlhorn, 2008). Wood and Fildes (1940) proposed that the mechanism of action of sulphonomides is related to the ability of the drug to inhibit the synthesis of folic acid by coccidian parasites through analogous competition to PABA (p-amino-benzoic acid). Finally, decoquinat acts by arresting the development of sporozoites following their penetration of the gut epithelium (Taylor and Bartram, 2012), probably through the inhibition of mitochondrial respiration and electron transport in *Eimeria* parasites (Wang, 1975, 1976; Fry and Williams, 1984). All the anticoccidials tested here seem to act against intracellular stages of the parasites, so they should have no effect against oocysts. This assumption has not been documented in the literature as far as we know. However, as expected, no inhibitory effect on oocyst sporulation rate was shown for any of the commercial or pure/source anticoccidials tested in the present study.

In conclusion, in this study we provide preliminary work towards the development of an *in vitro* model to evaluate ACE in ruminant hosts, using BCEC for cell culture studies and investigating different stages of development. One weakness of our study was the assumption that the anticoccidial concentrations remained stable when continuously included in the cell-culture medium throughout the study; however, we did not perform the necessary analyses to show that this was the case. We have been unable to determine the reason why, in general, we could not identify measurable effects from the different anticoccidials. Thus, further experiments, including the analysis of different cell lines, the implementation of cell permeability for non-ionophores anticoccidials, and investigation of their specific mechanisms of action are recommended. Despite these initial experiments not yielding any definitive clues, it is clear that the development of a suitable *in vitro* system for the evaluation of the ACE in ruminants would decrease the need for

animal experiments and could be used in the initial assessment of ACE of new anticoccidial drugs or bioactive substances. In our opinion, it therefore remains an important and worthy goal, and further investigations should be encouraged.

## Acknowledgements

This work was supported by The Foundation for Research Levy on Agricultural Products/Agricultural Agreement Research Fund, grant no. 244162 and Animalia (Norwegian Meat and Poultry Research Centre). We thank Esther Díaz and Berit Blomstrand for their excellent technical support. Finally, we cordially acknowledge Prof. Taubert and Prof. Hermosilla (Institute of Parasitology, Justus Liebig University of Giessen) who kindly donated the BCEC.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.exppara.2019.04.009>.

## Declaration of interest

The authors have no interests to declare.

## References

- Al-Rofaai, A., Rahman, W.A., Sulaiman, S.F., Yahaya, Z.S., 2012. In vitro activity of neem (*Azadirachta indica*) and cassava (*Manihot esculenta*) on three pre-parasitic stages of susceptible and resistant strains of *Teladorsagia (Ostertagia) circumcincta*. *Vet. Parasitol.* 188, 85–92.
- Alawa, C.B., Adamu, A.M., Gefu, J.O., Ajanusi, O.J., Abdu, P.A., Chiezey, N.P., Alawa, J.N., Bowman, D.D., 2003. In vitro screening of two Nigerian medicinal plants (*Vernonia amygdalina* and *Annona senegalensis*) for anthelmintic activity. *Vet. Parasitol.* 113, 73–81.
- Almeida, G.D., Feliz, D.C., Heckler, R.P., Borges, D.G., Onizuka, M.K., Tavares, L.E., Paiva, F., Borges, F.A., 2013. Ivermectin and moxidectin resistance characterization by larval migration inhibition test in field isolates of *Cooperia* spp. in beef cattle, Mato Grosso do Sul, Brazil. *Vet. Parasitol.* 191, 59–65. <https://doi.org/10.1016/j.vetpar.2012.08.012>.
- Araújo, S.A., Soares, A.M.D.S., Silva, C.R., Almeida Júnior, E.B., Rocha, C.Q., Ferreira, A.T.D.S., Perales, J., Costa-Júnior, L.M., 2017. In vitro anthelmintic effects of *Spigelia anthelmia* protein fractions against *Haemonchus contortus*. *PLoS One* 12, e0189803. <https://doi.org/10.1371/journal.pone.0189803>.
- Bach, U., Kalthoff, V., Mundt, H.C., Popp, A., Rinke, M., Dausgshies, A., Lüttge, B., 2003. Parasitological and morphological findings in porcine isosporosis after treatment with symmetrical triazinones. *Parasitol. Res.* 91, 27–33.
- Burke, J.M., Miller, J.E., Terrill, T.H., Orlik, S.T., Acharya, M., Garza, J.J., Mosjidis, J.A., 2013. *Sericea lespedeza* as an aid in the control of *Eimeria* spp. in lambs. *Vet. Parasitol.* 193, 3–46.
- Burt, S.A., Tersteeg-Zijdeveld, M.H., Jongerius-Gortemaker, B.G., Vervelde, L., Vernooij, J.C., 2013. In vitro inhibition of *Eimeria tenella* invasion of epithelial cells by phytochemicals. *Vet. Parasitol.* 191, 374–378. <https://doi.org/10.1016/j.vetpar.2012.09.001>.
- Chartier, C., Paraud, C., 2012. Coccidiosis due to *Eimeria* in sheep and goat, a review. *Small Rumin. Res.* 103, 84–92. <https://doi.org/10.1016/j.smallrumres.2011.10.022>.
- Chen, Y., Tang, Y., Guo, C., Wang, J., Boral, D., Nie, D., 2012. Nuclear receptors in the multidrug resistance through the regulation of drug-metabolizing enzymes and drug transporters. *Biochem. Pharmacol.* 83, 1112–1126. <https://doi.org/10.1016/j.bcp.2012.01.030>.
- Coles, G.C., Bauer, C., Borgsteede, F.H., Geerts, S., Klei, T.R., Taylor, M.A., Waller, P.J., 1992. World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) methods for the detection of anthelmintic resistance in nematodes of veterinary importance. *Vet. Parasitol.* 44, 35–44.
- Dausgshies, A., Najdrowski, M., 2005. Eimeriosis in cattle: current understanding. *J. Vet. Med. B Infect. Dis. Vet. Public. Health.* 52, 417–427.
- Deplazes, P., Eckert, J., Mathis, A., von Samson-Himmelstjerna, G., Zahner, H., 2016. Order eimeriida. In: *Parasitology in Veterinary Medicine*. Wageningen Academic Publishers, Wageningen, Netherlands, pp. 82–122.
- Diaferia, M., Veronesi, F., Morganti, G., Nisoli, L., Fioretti, D.P., 2013. Efficacy of toltrazuril 5% suspension (Baycox®, Bayer) and diclazuril (Vecoxan®, Janssen-Cilag) in the control of *Eimeria* spp. in lambs. *Parasitol. Res.* 112 (Suppl. 1), 163–168. <https://doi.org/10.1007/s00436-013-3440-1>.
- Enemark, H.L., Dahl, J., Enemark, J.M.D., 2015. Significance of timing on effect of methaphylactic toltrazuril treatment against eimeriosis in calves. *Parasitol. Res.* 114, S195–S206. <https://doi.org/10.1007/s00436-015-4526-8>.
- Fayer, R., Hammond, D.M., 1967. Development of first-generation schizonts of *Eimeria bovis* in cultured bovine cells. *J. Protozool.* 14, 764–772.
- Föllmann, W., Weber, S., Birkner, S., 2000. Primary cell cultures of bovine colon epithelium: isolation and cell culture of colonocytes. *Toxicol. Vitro* 14, 435–445.
- Foreyt, W.J., Hancock, D., Wescott, R.B., 1986. Prevention and control of coccidiosis in goats with decoquinate. *Am. J. Vet. Res.* 47, 333–335.
- Fry, M., Williams, R.B., 1984. Effects of decoquinate and clodipol on electron transport in mitochondria of *Eimeria tenella* (Apicomplexa: coccidia). *Biochem. Pharmacol.* 33, 229–240.
- Gadelhaq, S.M., Arafa, W.M., Abolhadid, S.M., 2018. In vitro activity of natural and chemical products on sporulation of *Eimeria* species oocysts of chickens. *Vet. Parasitol.* 251, 12–16.
- Gibbons, P., Love, D., Craig, T., Budke, C., 2016. Efficacy of treatment of elevated coccidial oocyst counts in goats using amprolium versus ponazuril. *Vet. Parasitol.* 218, 1–4.
- Harder, A., Haberkorn, A., 1989. Possible mode of action of toltrazuril: studies on two *Eimeria* species and mammalian and *Ascaris suum* enzymes. *Parasitol. Res.* 76, 8–12.
- Iqbal, Z., Lateef, M., Khan, M.N., Jabbar, A., Akhtar, M.S., 2006. Anthelmintic activity of *Sweritia chirata* against gastrointestinal nematodes of sheep. *Pfitoterapia* 77, 463–465.
- Iqbal, A., Tariq, K.A., Wazir, V.S., Singh, R., 2013. Antiparasitic efficacy of *Artemisia absinthium*, toltrazuril and amprolium against intestinal coccidiosis in goats. *J. Parasit. Dis.* 37, 88–93.
- Jackson, A.R., 1964. The isolation of viable coccidial sporozoites. *Parasitology* 54, 87–93.
- Jasso Díaz, G., Hernández, G.T., Zamilpa, A., Becerril Pérez, C.M., Ramírez Bribiesca, J.E., Hernández Mendo, O., Sánchez Arroyo, H., González Cortazar, M., Mendoza de Gives, P., 2017. In vitro assessment of *Argemone mexicana*, *Taraxacum officinale*, *Ruta chalepensis* and *Tagetes filifolia* against *Haemonchus contortus* nematode eggs and infective (L<sub>3</sub>) larvae. *Microb. Pathog.* 109, 162–168.
- Joachim, A., Altreuther, G., Bangoura, B., Charles, S., Dausgshies, A., Hinney, B., Lindsay, D.S., Mundt, H.C., Ocak, M., Sotiraki, S., 2018. W A A V P guideline for evaluating the efficacy of anticoccidials in mammals (pigs, dogs, cattle, sheep). *Vet. Parasitol.* 253, 102–119. <https://doi.org/10.1016/j.vetpar.2018.02.029>.
- Keeton, S.T.N., Navarre, C.B., 2018. Coccidiosis in large and small ruminants. *Vet. Clin. North. Am. Food Anim. Pract.* 34, 201–208.
- Khalafalla, R.E., Müller, U., Shahiduzzaman, M., Dyachenko, V., Desouky, A.Y., Alber, G., Dausgshies, A., 2011. Effects of curcumin (diferuloylmethane) on *Eimeria tenella* sporozoites in vitro. *Parasitol. Res.* 108, 879–886.
- Lan, L.H., Sun, B.B., Zuo, B.X.Z., Chen, X.Q., Du, A.F., 2017. Prevalence and drug resistance of avian *Eimeria* species in broiler chicken farms of Zhejiang province, China. *Poultry Sci.* 96, 2104–2109.
- Lanusse, C.E., Gascon, L.H., Prichard, R.K., 1995. Comparative plasma disposition kinetics of albendazole, fenbendazole and their metabolites in adult sheep. *J. Vet. Pharmacol. Ther.* 18, 196–203.
- Lim, J.H., Kim, M.S., Hwang, Y.H., Song, I.B., Park, B.K., Yun, H.I., 2010. Pharmacokinetics of toltrazuril and its metabolites, toltrazuril sulfoxide and toltrazuril sulfone, after a single oral administration to pigs. *J. Vet. Med. Sci.* 72, 1085–1087.
- Martínez-Valladares, M., Martínez-Pérez, J.M., Robles-Pérez, D., Cordero-Pérez, C., Famularo, M.R., Fernández-Pato, N., Castañón-Ordóñez, L., Rojo-Vázquez, F.A., 2013. The present status of anthelmintic resistance in gastrointestinal nematode infections of sheep in the northwest of Spain by *in vivo* and *in vitro* techniques. *Vet. Parasitol.* 191, 177–181.
- Matos, L., Muñoz, M.C., Molina, J.M., Ferrer, O., Rodríguez, F., Pérez, D., López, A.M., Martín, S., Hermosilla, C., Taubert, A., Ruiz, A., 2017a. Humoral immune responses of experimentally *Eimeria ninakohlyakimovae*-infected goat kids. *Comp. Immunol. Microbiol. Infect. Dis.* 51, 60–65. <https://doi.org/10.1016/j.cimid.2017.04.002>.
- Matos, L., Muñoz, M.C., Molina, J.M., Rodríguez, F., Pérez, D., López, A.M., Ferrer, O., Hermosilla, C., Taubert, A., Ruiz, A., 2017b. Protective immune responses during prepatency in goat kids experimentally infected with *Eimeria ninakohlyakimovae*. *Vet. Parasitol.* 242, 1–9. <https://doi.org/10.1016/j.vetpar.2017.04.016>.
- Matos, L., Muñoz, M.C., Molina, J.M., Rodríguez, F., Pérez, D., López, A.M., Hermosilla, C., Taubert, A., Ruiz, A., 2018. Age-related immune response to experimental infection with *Eimeria ninakohlyakimovae* in goat kids. *Res. Vet. Sci.* 118, 155–163. <https://doi.org/10.1016/j.rvsc.2018.02.004>.
- McDougal, L.R., Da Silva, J.M., Solis, J., Braga, M., 1987. A survey of sensitivity to anticoccidial drugs in 60 isolates of coccidia from broiler chickens in Brazil and Argentina. *Avian Dis.* 31, 287–292.
- Mehlhorn, H., 2008. DNA-synthesis-affecting drugs V: interference with dihydroorotate-dehydrogenase. In: Mehlhorn, H. (Ed.), *Encyclopedia of Parasitology*. Springer, Berlin, Heidelberg, pp. 389–392.
- Milhes, M., Guillerm, M., Robin, M., Eichstadt, M., Roy, C., Grisez, C., Prévot, F., Liénard, E., Bouhsira, E., Franc, M., Jacquet, P., 2017. A real-time PCR approach to identify anthelmintic-resistant nematodes in sheep farms. *Parasitol. Res.* 116, 909–920. <https://doi.org/10.1007/s00436-016-5364-z>.
- Molan, A.L., Liu, Z., De, S., 2009. Effect of pine bark (*Pinus radiata*) extracts on sporulation of coccidian oocysts. *Folia Parasitol.* 56, 1–5.
- Mundt, H.C., Bangoura, B., Rinke, M., Rosenbusch, M., Dausgshies, A., 2005. Pathology and treatment of *Eimeria zuernii* coccidiosis in calves: investigations in an infection model. *Parasitol. Int.* 54, 223–230.
- Novobilský, A., Stringano, E., Hayot Carbonero, C., Smith, L.M., Enemark, H.L., Mueller-Harvey, I., Thamsborg, S.M., 2013. In vitro effects of extracts and purified tannins of sainfoin (*Onobrychis vicifolia*) against two cattle nematodes. *Vet. Parasitol.* 196, 532–537. <https://doi.org/10.1016/j.vetpar.2013.03.024>.
- Odden, A., Enemark, H.L., Robertson, L.J., Ruiz, A., Hektoen, L., Stuen, S., 2017. Treatment against coccidiosis in Norwegian lambs and potential risk factors for development of anticoccidial resistance—a questionnaire-based study. *Parasitol. Res.* 116, 1237–1245. <https://doi.org/10.1007/s00436-017-5400-7>.
- Odden, A., Enemark, H.L., Ruiz, A., Robertson, L.J., Ersdal, C., Nes, S.K., Tømmerberg, V., Stuen, S., 2018a. Controlled efficacy trial confirming toltrazuril resistance in a field

- isolate of ovine *Eimeria* spp. Parasites Vectors 11, 394. <https://doi.org/10.1186/s13071-018-2976-4>.
- Odden, A., Denwood, M.J., Stuen, S., Robertson, L.J., Ruiz, A., Hamnes, I.S., Hektoen, L., Enemark, H.L., 2018b. Field evaluation of anticoccidial efficacy: a novel approach demonstrates reduced efficacy of toltrazuril against ovine *Eimeria* spp. in Norway. Int. J. Parasitol. Drugs Drug Resist. 8, 304–311. <https://doi.org/10.1016/j.ijpddr.2018.05.002>.
- Peek, H.W., Landman, W.J., 2005. Resistance of anticoccidial drugs of Dutch avian *Eimeria* spp. Field isolates originating from 1996, 1999 and 2001. Avian Pathol. 32, 391–401.
- Pérez, D., Ruiz, A., Muñoz, M.C., Molina, J.M., Hermosilla, C., López, A.M., Matos, L., Ortega, L., Martín, S., Taubert, A., 2015. Modulation of the pro-inflammatory molecules E-selectin and TNF- $\alpha$  gene transcription in *Eimeria ninakohlyakimovae*-infected primary caprine host endothelial cells. Parasitol. Int. 64, 471–477. <https://doi.org/10.1016/j.parint.2015.05.006>.
- Pérez, D., Muñoz, M.C., Molina, J.M., Muñoz-Caro, T., Silva, L.M., Taubert, A., Hermosilla, C., Ruiz, A., 2016. *Eimeria ninakohlyakimovae* induces NADPH oxidase-dependent monocyte extracellular trap formation and upregulates IL-12 and TNF- $\alpha$ , IL-6 and CCL2 gene transcription. Vet. Parasitol. 227, 143–150. <https://doi.org/10.1016/j.vetpar.2016.07.028>.
- R Core Team, 2017. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org>.
- Ramünke, S., Melville, L., Rinaldi, L., Hertzberg, H., de Waal, T., von Samson-Himmelstjerna, G., Cringoli, G., Mavrot, F., Skuce, P., Krücken, J., Demeler, J., 2016. Benzimidazole resistance survey for *Haemonchus*, *Teladorsagia* and *Trichostrongylus* in three European countries using pyrosequencing including the development of new assays for *Trichostrongylus*. Int. J. Parasitol. Drugs Drug Resist. 6, 230–240. <https://doi.org/10.1016/j.ijpddr.2016.10.002>.
- Rialch, A., Vatsya, S., Kumar, R.R., 2013. Detection of benzimidazole resistance in gastrointestinal nematodes of sheep and goats of sub-Himalayan region of northern India using different tests. Vet. Parasitol. 198, 312–318. <https://doi.org/10.1016/j.vetpar.2013.09.018>.
- Ruiz, A., González, J.F., Rodríguez, E., Martín, S., Hernández, Y.I., Almeida, R., Molina, J.M., 2006. Influence of climatic and management factors on *Eimeria* infections in goats from semi-arid zones. J. Vet. Med. B Infect. Dis. Vet. Public Health 53, 399–402.
- Ruiz, A., Behrendt, J., Zahner, H., Hermosilla, C., Pérez, D., Matos, L., Muñoz, M.C., Molina, J.M., Taubert, A., 2010. Development of *Eimeria ninakohlyakimovae* in vitro in primary and permanent cell lines. Vet. Parasitol. 173, 2–10. <https://doi.org/10.1016/j.vetpar.2010.05.023>.
- Ruiz, A., Guedes, A.C., Muñoz, M.C., Molina, J.M., Hermosilla, C., Martín, S., Hernández, Y.I., Hernández, A., Pérez, D., Matos, L., López, A.M., Taubert, A., 2012. Control strategies using diclazuril against coccidiosis in goat kids. Parasitol. Res. 110, 2131–2136. <https://doi.org/10.1007/s00436-011-2746-0>.
- Ruiz, A., Matos, L., Muñoz, M.C., Hermosilla, C., Molina, J.M., Andrada, M., Rodríguez, F., Pérez, D., López, A., Guedes, A.C., Taubert, A., 2013a. Isolation of an *Eimeria ninakohlyakimovae* field strain (Canary Islands) and analysis of its infection characteristics in goat kids. Res. Vet. Sci. 94, 277–284. <https://doi.org/10.1016/j.rvsc.2012.08.003>.
- Ruiz, A., Muñoz, M.C., Molina, J.M., Hermosilla, C., Rodríguez, F., Andrada, M., Martín, S., Guedes, A., Pérez, D., Matos, L., López, A.M., Taubert, A., 2013b. Primary infection of goats with *Eimeria ninakohlyakimovae* does not provide protective immunity against high challenge infections. Small Rumin. Res. 113, 258–266.
- Ruiz, A., Muñoz, M.C., Molina, J.M., Hermosilla, C., Andrada, M., Lara, P., Bordón, E., Pérez, D., López, A.M., Matos, L., Guedes, A.C., Falcón, S., Falcón, Y., Martín, S., Taubert, A., 2014. Immunization with *Eimeria ninakohlyakimovae*-live attenuated oocysts protect goat kids from clinical coccidiosis. Vet. Parasitol. 199, 8–17. <https://doi.org/10.1016/j.vetpar.2013.09.032>.
- Shresta, A., Freudenschuss, B., Jansen, R., Hinney, B., Ruttkowski, B., Joachim, A., 2017. Experimentally confirmed toltrazuril resistance in a field isolate of *Cystoisospora* suis. Parasites Vectors 10, 317. <https://doi.org/10.1186/s13071-017-2257-7>.
- Taylor, M.A., Bartram, D.J., 2012. The history of decoquinate in the control of coccidial infections in ruminants. J. Vet. Pharmacol. Ther. 35, 417–427. <https://doi.org/10.1111/j.1365-2885.2012.01421.x>.
- Taylor, M.A., Hunt, K.R., Goodyear, K.L., 2002. Anthelmintic resistance detection methods. Vet. Parasitol. 103, 183–194.
- Taylor, M.A., Marshall, R.N., Marshall, J.A., Catchpole, J., Bartram, D., 2011. Dose-response effects of diclazuril against pathogenic species of ovine coccidia and the development of protective immunity. Vet. Parasitol. 178, 48–57. <https://doi.org/10.1016/j.vetpar.2010.12.024>.
- Thabet, A., Alnassan, A.A., Dausgchies, A., Bangoura, B., 2015. Combination of cell culture and qPCR to assess the efficacy of different anticoccidials on *Eimeria tenella* sporozoites. Parasitol. Res. 114, 2155–2163. <https://doi.org/10.1007/s00436-015-4404-4>.
- Thabet, A., Zhang, R., Alnassan, A.A., Dausgchies, A., Bangoura, B., 2017. Anticoccidial efficacy testing: in vitro *Eimeria tenella* assays as replacement for animal experiments. Vet. Parasitol. 233, 86–96. <https://doi.org/10.1016/j.vetpar.2016.12.005>.
- Wang, C.C., 1975. Studies of the mitochondria from *Eimeria tenella* and inhibition of electron transport by coccidiostats. Biochem. Pharmacol. 25, 343–349.
- Wang, C.C., 1976. Inhibition of the respiration of *Eimeria tenella* by quinolone coccidiostats. Biochim. Biophys. Acta 396, 210–219.
- Woods, D.D., Fildes, P., 1940. The anti-sulphanilamide activity (*in vitro*) of paminobenzoic acid and related compounds. Chem. Ind. 59, 133–134.